


Jigar Sitapara, MD
Prime Facial Plastic Surgery
10285 Little Patuxent Parkway
Suite 250
301.302.7272

BROW LIFT POST-OPERATIVE INSTRUCTIONS

AMAZON POST-OP LIST: <https://a.co/cfYUsv8>

- Dr. Sitapara has curated these items to help you breeze through your recovery (*please review ingredients for any allergies or contraindications*):
 - L-Arginine Pro – Scientifically proven to aid in healing time and pain. Consume 1 scoop per day for the week before surgery and the week after surgery.
 - Vitamedica Bromelain – Studies show significant reduction in bruising. Take one pill per day for one week before surgery and the week after surgery.
 - Biocorneum Silicone Scar Gel – For use one week after surgery. Do not use on eyelid incisions.
 - General Gauze Sponges – Use as needed for wound care. Non-woven for comfort.
 - Extra Strength Tylenol – Lay a foundation of pain control before using additional pain medications for spikes in discomfort. 500mg rapid release tablets to take every 4-6 hours. Do not exceed 4000mg per day (8 tablets).
 - Active Skin Repair Hypochlorous Acid Hydrogel – Apply to incisions twice daily with clean hands. Allow it to dry before applying before bacitracin or other ointment. Do not apply to eye incisions.
 - Refresh Eye Drops – Sooth dry or irritated eyes as needed. Put the box in the fridge for added comfort.
 - Chloroseptic Spray – To soothe any throat irritation after surgery.
 - Sleep options - We generally recommend sleeping with your head slightly elevated after surgery to reduce swelling. We also want you to be strictly sleeping on your BACK. Explore these options and see which one works best for you.
 - AM Aeromax Back Sleep Training Pillow – Design trains you to sleep on your back. Recommend placing on top of another flat pillow to keep head elevated. Machine washable cover.
 - Cozymaker Wedge – Elevate your head and neck to an optimum angle. May need to combine with an airplane neck pillow to prevent side sleeping.
 - Airplane Pillow – *Hair Transplant Patients* Bring with you ON THE DAY OF SURGERY.

The following instructions apply to patients who have undergone brow lift. Since no two patients are ever exactly alike in their surgical needs, type of surgery performed or rate of healing, we may elect to individualize the following guidelines for each patient. In such instances, we will so instruct you. Otherwise, we urge you to follow the advice below very carefully, in order to accelerate your healing and maximize your surgical outcome.



Jigar Sitapara, MD
Prime Facial Plastic Surgery
10285 Little Patuxent Parkway
Suite 250
301.302.7272

Preoperative Instructions

If possible, do not take any Aspirin or NSAID medications (ibuprofen, motrin, advil, etc) for 14 days prior to your procedure, and then only on the advice of your personal physician. Please let your surgeon know if this is not possible prior to your surgery.

Avoiding alcohol for 7 days prior to surgery can help with the healing process.

Postoperative Instructions

You may be up and around the day after surgery, but some natural fatigue may persist for 2-3 days due to the normal effects of the anesthesia and surgical procedure.

When you move, stand or change positions, do so deliberately and carefully for the first 7 days. Be careful not to pull down on your face.

You may eat a normal diet the day following the surgery. In moderation, talking and smiling are perfectly acceptable. It is normal to have tightness when opening your mouth. This will resolve in time.

Your head should be elevated on at least two pillows during sleep for the first 14 days, in order to keep your head higher than your heart to help facilitate the resolution of swelling. Do not sleep face down, rather on your back or side.


DO NOT TAKE ANY ASPIRIN OR ASPIRIN CONTAINING MEDICINES for 14 days prior to and after the surgery, and then only on the advice of your personal physician. Other routinely taken medications may be taken as necessary.

Any unexplained development of pain, facial swelling or fever should be reported to us immediately.

Some facial swelling and bruising are normally present after brow lift, but the degree of each varies widely from patient to patient. Do not be concerned if you have more or less than others who have undergone the "same" operation. Generally, most patients appear quite sociably acceptable within 10-14 days after surgery.

You may gently cleanse the incision lines twice daily with the HYPOCHLOROUS ACID GEL (or hydrogen peroxide if you did not purchase this item). Apply bacitracin provided sparingly twice daily to the incision lines in order to avoid excessive crusting of the incisions and to accelerate the reduction of incision redness. Do not apply any other ointment or medications unless we prescribe it.

You may gently shampoo your hair 72 hours after surgery, avoiding any strong rubbing or combing trauma to the incisions in the hair and around the ear. Do not blow dry for 5 days, and postpone any planned permanent waves or hair coloring for 4 weeks following surgery.



Jigar Sitapara, MD
Prime Facial Plastic Surgery
10285 Little Patuxent Parkway
Suite 250
301.302.7272

Excessive exposure to sun (including sun-tanning parlors) in the first 3 weeks after surgery may result in prolonged facial swelling and injury to the skin. Thereafter, you should always protect your skin with a sun-screen in order to decrease the inevitable aging effects of the sun on your skin.

Makeup may be worn 48 hours after surgery.

We greatly appreciate the confidence you have shown in us by allowing us to assist you in improving your appearance and health, and you may be assured of our very best efforts to achieve the best surgical result possible for your particular individual anatomy and condition.